

ZBW-Ausgabe 8-9/2015

Literaturverzeichnis zum Fortbildungsbeitrag

Aplasia der oberen seitlichen Schneidezähne und multiple Nichtanlagen (Teil 2) Therapeutische Möglichkeiten

**Prof. Dr. Dr. Bernd Lapatki Dr. Johanna Radeke Klinik für Kieferorthopädie und
Orthodontie, Universitätsklinikum Ulm**

Referenzen

1. Yin, W. and Z. Bian, The Gene Network Underlying Hypodontia. *J Dent Res*, 2015.
2. Mattheeuws, N., L. Dermaut, and G. Martens, Has hypodontia increased in Caucasians during the 20th century? A meta-analysis. *Eur J Orthod*, 2004. 26(1): p. 99-103.
3. Polder, B.J., et al., A meta-analysis of the prevalence of dental agenesis of permanent teeth. *Community Dent Oral Epidemiol*, 2004. 32(3): p. 217-26.
4. Kim, Y.H., Investigation of hypodontia as clinically related dental anomaly: prevalence and characteristics. *ISRN Dent*, 2011. 2011: p. 246135.
5. Rakhshan, V., Congenitally missing teeth (hypodontia): A review of the literature concerning the etiology, prevalence, risk factors, patterns and treatment. *Dent Res J (Isfahan)*, 2015. 12(1): p. 1-13.
6. Kepes, D. and K. Gabris, [Prevalence of hypodontia at the department of pedodontics and orthodontics of Semmelweis University]. *Fogorv Sz*, 2013. 106(1): p. 23-6.
7. Friedrich, R.E., A. Leist, and H.A. Scheuer, [Eruption times of permanent teeth in children and adolescents in the German state of Saarland]. *Arch Kriminol*, 2008. 222(3-4): p. 73-104.
8. Kochhar, R. and A. Richardson, The chronology and sequence of eruption of human permanent teeth in Northern Ireland. *Int J Paediatr Dent*, 1998. 8(4): p. 243-52.
9. Moorrees, C.F., E.A. Fanning, and E.E. Hunt, Jr., AGE VARIATION OF FORMATION STAGES FOR TEN PERMANENT TEETH. *J Dent Res*, 1963. 42: p. 1490-502.
10. Bicakci, A.A., C. Doruk, and H. Babacan, Late development of a mandibular second premolar. *Korean J Orthod*, 2012. 42(2): p. 94-8.
11. Moraschini, V., et al., Evaluation of survival and success rates of dental implants reported in longitudinal studies with a follow-up period of at least 10 years: a systematic review. *Int J Oral Maxillofac Surg*, 2015. 44(3): p. 377-88.
12. Setzer, F.C. and S. Kim, Comparison of long-term survival of implants and endodontically treated teeth. *J Dent Res*, 2014. 93(1): p. 19-26.
13. Northway, W., Hemisection: one large step toward management of congenitally missing lower second premolars. *Angle Orthod*, 2004. 74(6): p. 792-9.
14. Valencia, R., M. Saadia, and G. Grinberg, Controlled slicing in the management of congenitally missing second premolars. *Am J Orthod Dentofacial Orthop*, 2004. 125(5): p. 537-43.
15. Wilmes, B., et al., Mini-implant-supported temporary pontics. *J Clin Orthod*, 2014. 48(7): p. 422-9.
16. Bjerklin, K. and J. Bennett, The long-term survival of lower second primary molars in subjects with agenesis of the premolars. *Eur J Orthod*, 2000. 22(3): p. 245-55.
17. Nance, H.N., The limitations of orthodontic treatment; mixed dentition diagnosis and treatment. *Am J Orthod*, 1947. 33(4): p. 177-223.